

Approximation Methods for Nonlinear m -Accretive Operator Equations*

M. O. Osilike

Department of Mathematics, University of Nigeria, Nsukka, Nigeria

Submitted by John Horváth

Received March 12, 1996

Suppose E is a real reflexive Banach space and $T : D(T) \subset E \rightarrow E$ is a locally Lipschitzian m -accretive operator, where the domain of T , $D(T)$, is a proper subset of E . For any $f \in E$, approximation methods are constructed which converge strongly to the solution of the equation $x + Tx = f$. Explicit error estimates are given and convergence is at least as fast as a geometric progression. © 1997 Academic Press

INTRODUCTION

Let E be a real Banach space.

An operator T with domain $D(T)$ and range $R(T)$ in E is called *accretive* if the inequality

$$\|x - y\| \leq \|x - y + \lambda(Tx - Ty)\| \quad (1)$$

holds for all $x, y \in D(T)$ and for all $\lambda > 0$. If T is accretive and $(I + rT)(D(T)) = E$ for all $r > 0$ then T is called *m -accretive*, where I denotes the identity operator.

If T is m -accretive, then for any given $f \in E$ the equation

$$x + Tx = f \quad (2)$$

has a unique solution.

Recently, Chidume and the author [2] and Liang [4] studied methods for approximating the solution of Eq. (2) in *uniformly smooth Banach spaces*

*Research supported by a grant from TWAS (94-224 RG/MATHS/AF/AC), ICTP, Trieste, Italy.

under the natural setting that the domain of T , $D(T)$, is a proper subset of E , T maps $D(T)$ into E , and is m -accretive. In this note we continue the study of methods of approximating the solution of Eq. (2) (when T is m -accretive) in Banach spaces much more general than those considered in Chidume and the author [2] and Liang [4] under the natural setting that the domain of T , $D(T)$, is a proper subset of E and T maps $D(T)$ into E . In particular, we extend Theorems 1 and 2 of Liang [4] from *real uniformly smooth Banach spaces* to the much more general *real reflexive Banach spaces*.

1. MAIN RESULTS

In the sequel L will denote the local Lipschitz constant of T .

THEOREM 1. *Suppose E is a real reflexive Banach space, and $T : D(T) \subset E \rightarrow E$ is m -accretive and locally Lipschitzian. Suppose $D(T)$ is open and that $x^* \in D(T)$ is the unique solution of the equation $x + Tx = f$, $f \in E$. Suppose $\{\alpha_n\}_{n=0}^\infty$ is a real sequence satisfying the conditions:*

- (i) $0 \leq \alpha_n \leq 1/2[L^2 + 2L + 2]$, $n \geq 0$
- (ii) $\sum_{n=0}^\infty \alpha_n = \infty$.

Then there exists a closed convex neighborhood B of x^ contained in $D(T)$ and for any given $x_0 \in B$, a sequence $\{x_n\}_{n=0}^\infty$ of elements of B such that on setting*

$$p_n = (1 - \alpha_n)x_n + \alpha_n(f - Tx_n), \quad n \geq 0$$

the sequence $\{p_n\}$ satisfies the condition

$$\|p_{n-1} - x_n\| = \inf\{\|p_{n-1} - x\| : x \in B\} \quad \forall n \geq 1 \quad (3)$$

and converges strongly to x^ . Moreover, if $\alpha_n = 1/2[L^2 + 2L + 2]$ for all $n \geq 0$ then*

$$\|p_n - x^*\| \leq \rho^n \|p_0 - x^*\|,$$

where $\rho = (1 - 1/4[L^2 + 2L + 2]) \in (0, 1)$.

Proof. Define $S : D(T) \rightarrow E$ by $Sx = f - Tx$. Observe that x^* is a fixed point of S and that S is locally Lipschitz with constant L . Furthermore $(-S)$ is accretive so that for all $r > 0$, and $x, y \in D(T)$

$$\|x - y\| \leq \|x - y - r(Sx - Sy)\|. \quad (4)$$

Without loss of generality we may assume $L \geq 1$. Let $B(y, r) = \{x \in E : \|x - y\| \leq r\}$. Then there exists $r_1 > 0$ such that $B(x^*, r_1) \subseteq D(T)$.

Since S is locally Lipschitzian, there exists $r_2 > 0$ such that S is Lipschitzian on $B(x^*, r_2)$. Let $r = \min\{r_1, r_2\}$. Then $B(x^*, r) \subseteq D(T)$ and S is Lipschitzian on $B(x^*, r)$. Let $B = B(x^*, r/2L)$. Given any $x_0 \in B$, then $\|Sx_0 - x^*\| \leq r/2 < r$ so that $Sx_0 \in B(x^*, r)$. Hence $p_0 = (1 - \alpha_0)x_0 + \alpha_0 Sx_0 \in B(x^*, r)$. Since E is reflexive, there exists $x_1 \in B$ such that

$$\|p_0 - x_1\| = \inf\{\|p_0 - x\| : x \in B\}.$$

Thus

$$p_1 = (1 - \alpha_1)x_1 + \alpha_1 Sx_1 \in B(x^*, r).$$

By continuing this process we obtain $\{p_n\}$ in $B(x^*, r)$ and $\{x_n\}$ in B satisfying the conditions

$$\begin{aligned} p_n &= (1 - \alpha_n)x_n + \alpha_n Sx_n, & n \geq 0 \\ \|p_{n-1} - x_n\| &= \inf\{\|p_{n-1} - x\| : x \in B\}, & n \geq 1. \end{aligned} \quad (5)$$

Thus

$$\|x_n - x^*\| \leq \|p_{n-1} - x^*\|, \quad n \geq 1.$$

We now prove that $\lim p_n = x^*$. From (5) we obtain

$$\begin{aligned} x_n &= p_n + \alpha_n x_n - \alpha_n Sx_n \\ &= (1 + \alpha_n)p_n - \alpha_n Sp_n + \alpha_n^2(x_n - Sx_n) + \alpha_n(Sp_n - Sx_n). \end{aligned} \quad (6)$$

Observe that

$$x^* = (1 + \alpha_n)x^* - \alpha_n Sx^*. \quad (7)$$

Thus from (6) and (7) we obtain

$$\begin{aligned} \|x_n - x^*\| &= \|(1 + \alpha_n)(p_n - x^*) - \alpha_n(Sp_n - Sx^*) \\ &\quad + \alpha_n^2(x_n - Sx_n) + \alpha_n(Sp_n - Sx_n)\| \\ &\geq (1 + \alpha_n)\|p_n - x^*\| - \frac{\alpha_n}{1 + \alpha_n}\|Sp_n - Sx^*\| \\ &\quad - \alpha_n^2\|x_n - Sx_n\| - \alpha_n\|Sp_n - Sx_n\| \\ &\geq (1 + \alpha_n)\|p_n - x^*\| - \alpha_n^2\|x_n - Sx_n\| - \alpha_n\|Sp_n - Sx_n\| \\ &\quad \text{(using (4)).} \end{aligned}$$

Hence

$$\begin{aligned}
 \|P_n - x^*\| &\leq \frac{1}{1 + \alpha_n} \|x_n - x^*\| + \alpha_n^2 \|x_n - Sx_n\| + \alpha_n \|Sp_n - Sx_n\| \\
 &\leq [1 - \alpha_n + \alpha_n^2] \|x_n - x^*\| \\
 &\quad + (1 + L) \alpha_n^2 \|x_n - x^*\| + L(1 + L) \alpha_n^2 \|x_n - x^*\| \\
 &= [1 - \alpha_n + (L^2 + 2L + 2) \alpha_n^2] \|x_n - x^*\| \\
 &\leq [1 - \frac{1}{2} \alpha_n] \|p_{n-1} - x^*\| \\
 &\leq \exp\left(-\frac{1}{2} \sum_{j=0}^n \alpha_j\right) \|p_0 - x^*\| \rightarrow 0 \quad \text{as } n \rightarrow \infty.
 \end{aligned} \tag{8}$$

If we set $\alpha_n = 1/2[L^2 + 2L + 2] \forall n \geq 0$ in (8) we obtain

$$\|p_n - x^*\| \leq \left[1 - \frac{1}{4(L^2 + 2L + 2)}\right] \|p_{n-1} - x^*\| \leq \rho^n \|p_0 - x^*\|,$$

completing the proof of Theorem 1.

Remark 1. In proving Theorem 1 we first observed that the accretivity of T implies that $(-S)$ is accretive and hence inequality (4) holds. In order to apply inequality (4) we used (5) to express x_n in the form (6). Furthermore, we expressed x^* in the form (7). Using (6) and (7) and an application of inequality (4) and the Lipschitz condition of S we were able to prove Theorem 1 without the application of an inequality of Reich used in [4] and which hold in uniformly smooth Banach spaces. Moreover, we did not apply a lemma of Weng which was also used in [4].

Remark 2. Theorem 1 extends Theorems 1 and 2 of Liang [4] from real uniformly smooth Banach spaces to real reflexive Banach spaces. Moreover, our theorem provides additional information about the error estimates. Convergence is at least as fast as a geometric progression.

Remark 3. Using straightforward modifications of the proof of Theorem 1 one can easily prove the following theorem for a more general iteration scheme.

THEOREM 2. Suppose $E, T, D(T), S$, and x^* are as in Theorem 1. Suppose $\{\alpha_n\}_{n=0}^\infty$ and $\{\beta_n\}_{n=0}^\infty$ are real sequences satisfying the conditions:

- (i) $0 \leq \alpha_n \leq 1/2[L^2 + 2L + 2], \quad n \geq 0$
- (ii) $0 \leq \beta_n \leq 1/4[L^2(1 + L)], \quad n \geq 0$
- (iii) $\sum_{n=0}^\infty \alpha_n = \infty.$

Then there exists a closed convex neighborhood B of x^* contained in $D(T)$ and for any given $x_0 \in B$, a sequence $\{x_n\}_{n=0}^{\infty}$ of elements of B such that on setting

$$y_n = (1 - \beta_n)x_n + \beta_n Sx_n, \quad n \geq 0$$

$$p_n = (1 - \alpha_n)x_n + \alpha_n Sy_n, \quad n \geq 0$$

the sequence $\{p_n\}$ satisfies the condition

$$\|p_{n-1} - x_n\| = \inf\{\|p_{n-1} - x\| : x \in B\} \quad \forall n \geq 1$$

and converges strongly to x^* . Moreover, if $\alpha_n = 1/2[L^2 + 2L + 2] \forall n \geq 0$, and $\beta_n = 1/4L^2(1 + L) \forall n \geq 0$ then

$$\|p_n - x^*\| \leq \rho^n \|p_0 - x^*\|,$$

where $\rho = (1 - 1/8[L^2 + 2L + 2]) \in (0, 1)$.

ACKNOWLEDGMENTS

The author is grateful to the referee for making useful comments and suggestions, and to Professor C. H. Morales for useful discussions at the National Mathematical Centre, Abuja, Nigeria.

REFERENCES

1. F. E. Browder, Nonlinear operators and nonlinear equations of evolution in Banach spaces, in "Proc. Sympos. Pure Math.," Vol. 18, Part 2, Amer. Math. Soc., Providence, Rhode Island, 1976.
2. C. E. Chidume and M. O. Osilike, Approximation methods for nonlinear operator equations of the m -accretive type, *J. Math. Anal. Appl.* **189** (1995), 225–239.
3. T. Kato, Nonlinear semigroups and evolution equations, *J. Math. Soc. Japan* **19** (1967), 508–520.
4. Z. Liang, Iterative solution of nonlinear equations involving m -accretive operators in Banach spaces, *J. Math. Anal. Appl.* **188** (1994), 410–416.